

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

U.S. Navy Research

U.S. Department of Defense

2019

First Report of New Delhi Metallo- β -Lactamase Carbapenemase–Producing *Acinetobacter baumannii* in Peru

Claudio Rocha

U.S. Naval Medical Research Unit No. 6, claudiorochacalderon@gmail.com

Manuela Bernal

U.S. Naval Medical Research Unit No. 6

Enrique Canal

U.S. Naval Medical Research Unit No. 6

Paul Rios

U.S. Naval Medical Research Unit No. 6

Rina Meza

U.S. Naval Medical Research Unit No. 6

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/usnavyresearch>

Rocha, Claudio; Bernal, Manuela; Canal, Enrique; Rios, Paul; Meza, Rina; Lopez, Miguel; Burga, Rosa; Abadie, Ricardo; Pizango, Melita; Diaz, Elia; Briones, Alexander; Ramal-Asayag, Cesar; Vicente, William; Regeimbal, James; and McCoy, Andrea, "First Report of New Delhi Metallo- β -Lactamase Carbapenemase–Producing *Acinetobacter baumannii* in Peru" (2019). *U.S. Navy Research*. 152. <https://digitalcommons.unl.edu/usnavyresearch/152>

This Article is brought to you for free and open access by the U.S. Department of Defense at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in U.S. Navy Research by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Claudio Rocha, Manuela Bernal, Enrique Canal, Paul Rios, Rina Meza, Miguel Lopez, Rosa Burga, Ricardo Abadie, Melita Pizango, Elia Diaz, Alexander Briones, Cesar Ramal-Asayag, William Vicente, James Regeimbal, and Andrea McCoy

First Report of New Delhi Metallo- β -Lactamase Carbapenemase–Producing *Acinetobacter baumannii* in Peru

Claudio Rocha,^{1*} Manuela Bernal,¹ Enrique Canal,¹ Paul Rios,¹ Rina Meza,¹ Miguel Lopez,¹ Rosa Burga,¹ Ricardo Abadie,¹
Melita Pizango,¹ Elia Diaz,² Alexander Briones,² Cesar Ramal-Asayag,^{2,3} William Vicente,⁴ James Regeimbal,^{1*}
and Andrea McCoy¹

¹U. S. Naval Medical Research Unit No 6, Lima, Peru; ²Hospital Regional de Loreto, Loreto, Peru; ³Universidad Nacional de la Amazonia Peruana, Loreto, Peru; ⁴Instituto Nacional de Enfermedades Neoplasicas, Lima, Peru

Abstract. Here we report the first incidence of New Delhi metallo- β -lactamase (NDM-1)–producing *Acinetobacter baumannii* in Peru, identified via a strain-based nosocomial surveillance project carried out in Lima and Iquitos. The *bla*_{NDM-1} gene was detected by multiplex polymerase chain reaction (PCR) and confirmed by loci sequencing. *Acinetobacter baumannii* is a nearly ubiquitous and promiscuous nosocomial pathogen, and the acquisition of *bla*_{NDM-1} by *A. baumannii* may facilitate an increase in the prevalence of this important resistance marker in other nosocomial pathogens.

INTRODUCTION

Health-care-associated infections (HAI) and antimicrobial resistance are of great public health concern worldwide.¹ Bacterial pathogens causing HAI have become increasingly more resistant over the past 10–15 years as a result of various mechanisms, including gene-mediated enzymes such as class A, B, and D β -lactamases.² Numerous factors, from the horizontal transfer of gene-encoded enzymes to global travel, have allowed resistance and resistant organisms to rapidly spread with great clinical and epidemiological impact.^{3,4} The recently described carbapenemase, New Delhi metallo- β -lactamase (NDM-1) class B, is of particular epidemiological and clinical concern. Since its discovery in 2008, NDM-1 has rapidly spread worldwide and confers resistance to almost all lactams, with the exception of aztreonam, leaving limited therapeutic options against pathogens harboring NDM-1, typically colistin, tigecycline, and fosfomycin.^{2,5,6} In Latin America, the *bla*_{NDM-1} gene was first reported in 2011 from Guatemala and Colombia, and later from Mexico in 2012, Brazil in 2013, and Uruguay in 2013, with all instances from Enterobacteriaceae.⁵ The *bla*_{NDM-1} gene in non-fermentative pathogens was first reported in Latin America from Honduras (*Acinetobacter baumannii*) and Paraguay (*Acinetobacter pittii*) in 2012, and later from Brazil (*A. baumannii*) in 2014, Cuba (*Acinetobacter soli*) in 2015, Argentina (*Acinetobacter junii*) in 2016, and Colombia (*A. baumannii*) in 2017.^{5–8} In Peru, the first report of *bla*_{NDM-1} was in May 2017 in a set of nine *Klebsiella pneumoniae* infecting or colonizing critically ill neurological patients from one hospital in Lima.⁹ Here, we describe the identification of the first three strains of *A. baumannii* harboring *bla*_{NDM-1} in Peru as part of a strain-based surveillance project carried out by the Naval Medical Research Unit No Six (NAMRU-6) in Lima, Peru.

MATERIALS AND METHODS

In June 2016, NAMRU-6 began ongoing strain collection from Lima (three hospitals) and Iquitos (two hospitals), Peru. Strain collection for this effort focuses on, but is not limited to,

multidrug-resistant or extremely drug-resistant (MDR/XDR) ESKAPE pathogens (*Enterococcus faecium*, *Staphylococcus aureus*, *K. pneumoniae*, *A. baumannii*, *Pseudomonas aeruginosa*, and *Enterobacter* spp.), described as the most important pathogens causing HAI globally.¹⁰

At participating hospitals in Lima and Iquitos, clinicians order sampling and cultures as part of routine patient care (hemoculture, wound swabs, secretion sampling, catheter tip cultures, etc.), and the local hospital laboratory or infection control program isolates and identifies the etiology of the infection, as well as performs susceptibility testing. Briefly, at both sites, conventional culture procedures are performed for identification and susceptibility testing using routine biochemical algorithms and disk diffusion tests, respectively. In addition, at the participating hospital in Lima, isolates from bacteremic patients were recovered using the automated BD BACTEC blood culture system and identification and susceptibilities were confirmed using a Phoenix system (Becton & Dickinson Diagnostics, Sparks, MD).

Once isolates are identified at the participating hospital laboratory, the strains are cryopreserved and transferred within 2 weeks to NAMRU-6 for confirmation and further molecular characterization. No clinical, epidemiological, or personally identifiable data are obtained by NAMRU-6; only limited data about how a strain was isolated are reported.

At NAMRU-6, all strains such as those reported here are confirmed by routine biochemical algorithms and matrix-assisted laser desorption/ionization-time-of-flight (MALDI-TOF) (Becton & Dickinson Bruker Microflex, Bremen, Germany). Antibiotic susceptibility testing (AST) is performed by using the disk diffusion test and the Phoenix system (BD Diagnostics, Sparks, MD), and results are interpreted using the Clinical Laboratory Standards Institute guidelines.¹¹ Molecular characterization of strains is performed on highly resistant MDR/XDR pathogens as described by Magiorakos et al.¹² Each MDR/XDR isolate is first characterized by multiplex PCR. Colony PCR and cycling conditions are performed essentially as described by Poirer et al.,¹³ using the following primers that amplify loci within each open reading frame as described previously^{13,14}:

IMP-F: 5'-GGAATAGAGTGGCTTAAYTCTC-3', IMP-R: 3'-GTTTAAAYAAAACAACCACC-5' (232 bp); VIM-F: 5'-GATGGTGTTTGGTCGCATA-3', VIM-R: 3'-CGAATGCGCAGCACAG-5' (390 bp); NDM-F: 5'-GGTTTGGCGATCTGGTTTTTC-3', NDM-R: 3'-CGGAATGGCTCATCACGATC-5' (621 bp);

* Address correspondence to Claudio Rocha, Universidad Autonoma de Barcelona, Barcelona, Spain, E-mail: claudiochacalderon@gmail.com or James Regeimbal, Hospital Naval, Av. Venezuela cuadra 36 S/N, Bellavista, Callao, Peru, E-mail: james.m.regeimbal.mil@mail.mil.

and KPC-F: 5'-CGTCTAGTTCTGCTGCTTG-3', KPC-R: 3'-CTTGTCATCCTTGTTAGGCG-5' (798 bp).

Cleaned colony PCR products from each PCR-positive isolate are then subjected to amplicon sequencing using an ABI 3130XL genetic analyzer (Applied Biosystems, New York, NY) according to the manufacturer's protocol, using the same primers listed previously. Sequence analysis is performed using Sequencher V5.4 Gene Codes Corporation (Ann Arbor, MI) and basic local alignment search tool (BLAST, National Center for Biotechnology Information) to compare obtained target sequences with the database. Thus, MDR/XDR strains are characterized by NAMRU-6 for phenotypic and genotypic resistance to the partial sequence level.

RESULTS

From June 2016 to November 2017, 875 ESKAPE and 213 *Escherichia coli* isolates were collected from Lima and Iquitos hospitals (Table 1). The most frequent pathogen collected from Lima was *P. aeruginosa* (36%), whereas that of Iquitos was *E. coli* (46%). Enterobacteriaceae (*K. pneumoniae*, *Enterobacter* spp., and *E. coli*) represented 45% of all pathogens collected, whereas *A. baumannii* represented 11%. From the total 1,088 strains collected, 978 were Gram-negative pathogens and AST was performed on 974. Of the 974 isolates, 472 (48%) were non-susceptible (resistant or intermediate phenotype) to carbapenem. PCR was performed on 764 (78%) of the 974 strains with AST results, and 155 (20%) of those were positive for at least one of the four carbapenemase-encoding genes tested. Interestingly, 8 (5%) strains harboring PCR-detectable carbapenemase-encoding genes were susceptible to carbapenems, indicating not only a general agreement between PCR and AST but also that there are pseudogenes for carbapenem resistance present in the susceptible clinical isolates here. Such sequences likely have an interesting role in the dynamics of resistance within bacterial populations, and further analysis of such sequences in populations of clinical isolates in Peru is needed. From these 155 strains, an additional eight non-susceptible strains harboring *bla*_{NDM-1} were detected: five Enterobacteriaceae (one *E. coli* and four *K. pneumoniae*, all from Lima) and three *A. baumannii* (one from Lima and two from Iquitos).

The only *A. baumannii* harboring *bla*_{NDM-1} from Lima was isolated in January 2017 from a hemoculture taken in a tertiary care oncology hospital and was classified as MDR using the same criteria as previously mentioned.¹² The strain was resistant to all β -lactams with the exception of aztreonam, to which it showed intermediate resistance. This strain was also resistant to all the carbapenems tested, as well as to

fosfomycin. In addition, the strain showed intermediate resistance to tetracycline but was sensitive to quinolones, aminoglycosides, trimethoprim-sulfamethoxazole, and colistin (Table 2). The two *A. baumannii* strains from Iquitos harboring *bla*_{NDM-1} were isolated from the same tertiary care general hospital but at different times and from different types of infections. The first was also isolated in January 2017 from a central venous catheter tip and was classified as MDR. This strain was resistant to all β -lactams with the exception of piperacillin-tazobactam, to which it showed intermediate resistance, and it was also resistant to all the carbapenems tested and fosfomycin. In addition, this strain also showed intermediate resistance to ciprofloxacin but was sensitive to tetracycline, trimethoprim-sulfamethoxazole, levofloxacin, aminoglycosides, and colistin. The second *A. baumannii* from Iquitos was isolated in February 2017 from a postsurgical wound and was classified as XDR. This strain was resistant to all the antibiotics tested except ampicillin-sulbactam, to which it showed intermediate resistance, and colistin and tetracycline, to which it was sensitive (Table 2). Amplicon sequencing of all three *A. baumannii* strains returned a 99% match to *bla*_{NDM-1}.

DISCUSSION

Carbapenem-resistant *A. baumannii* is a serious public health concern worldwide.^{2,6,15,16} *Acinetobacter baumannii* has spread throughout the world acquiring antibiotic resistance, first to the most common antibiotics (β -lactams, sulfonamides, and aminoglycosides) and later to carbapenems.^{15,17,18} The clinical

TABLE 2

Resistance pattern of the three strains of *Acinetobacter baumannii* harboring *bla*_{NDM-1} gene

Code	NSC2058	NSI1409	NSI1428
Study site	Lima	Iquitos	Iquitos
Sample type	Hemoculture	CVC tip	Wound
Sample date	January 27, 2017	January 18, 2017	January 17, 2017
AMK	S	S	R
GEN	S	S	R
AMP	R	R	R
SAM	R	R	I
TIC	R	R	R
TIM	R	R	R
TZP	R	I	R
ATM	I	R	R
CAZ	R	R	R
CFZ	R	R	R
CRO	R	R	R
FOX	R	R	R
FEP	R	R	R
LVX	S	S	R
CIP	S	I	R
IPM	R	R	R
MEM	R	R	R
FOF	R	R	R
SXT	S	S	R
TET	I	S	S
CST	S	S	S
Definition for acquired resistance	MDR	MDR	XDR

AMK = amikacin; AMP = ampicillin; ATM = aztreonam; CAZ = ceftazidime; CFZ = cefazolin; CIP = ciprofloxacin; CRO = ceftriaxone; CST = colistin; CVC = central venous catheter; FEP = cefepime; FOF = fosfomycin; FOX = ceftiofur; GEN = gentamicin; I = intermediate; IPM = imipenem; LVX = levofloxacin; MDR = multidrug resistant; MEM = meropenem; NDM-1 = New Delhi metallo- β -lactamase; R = resistant; S = sensitive; SAM = ampicillin-sulbactam; SXT = trimethoprim-sulfamethoxazole; TET = tetracycline; TIC = ticarcillin; TIM = ticarcillin-clavulanic acid; TZP = piperacillin-tazobactam; XDR = extremely drug resistant.

TABLE 1

ESKAPE strains and *Escherichia coli* collected from June 2016 to November 2017 from Lima and Iquitos, Peru

ESKAPE	Lima	Iquitos	Total
<i>Enterococcus faecium</i>	20	0	20 (1.8%)
<i>Enterobacter</i> spp.	38	2	40 (3.7%)
<i>Staphylococcus aureus</i>	80	10	90 (8.3%)
<i>Acinetobacter baumannii</i>	116	8	124 (11.4%)
<i>Klebsiella pneumoniae</i>	204	34	238 (21.9%)
<i>Pseudomonas aeruginosa</i>	343	20	363 (33.4%)
Total (ESKAPE)	801	74	875 (80.4%)
<i>E. coli</i>	150	63	213 (19.6%)
Total (ESKAPE and <i>E. coli</i>)	951	137	1,088

ESKAPE = *Enterobacter* spp., *S. aureus*, *K. pneumoniae*, *A. baumannii*, *P. aeruginosa*, and *E. faecium*.

impact of this pathogen stems from its ability to upregulate or readily acquire resistance determinants and survive for prolonged periods in nosocomial environments, thus enhancing its rapid dissemination.^{6,16,17} Carbapenem-resistant *A. baumannii* is mainly thought to produce OXA- β -lactamases (oxacillin-hydrolysing) and is less frequently an NDM-1 producer.^{15–17} Conversely, some studies have found truncated forms of the *bla*_{NDM-1}-harboring composite transposons Tn725 in Enterobacteriaceae species, whereas the complete sequence can be found in *A. baumannii*. Taken together, this suggests that *A. baumannii* may serve as a reservoir of *bla*_{NDM-1} that can be transferred to other pathogens.^{2,6,19,20} Furthermore, the spread of *A. baumannii* harboring *bla*_{NDM-1} is likely not a matter of a single lineage but may involve different contemporary clones.

NSC2058 and NSI1409 were found in Lima and Iquitos, respectively; yet they have similar patterns of resistance for the antibiotics tested (Table 2), and both were found in January 2017. Together these results suggest a possible epidemiological link, despite the fact that Iquitos can only be reached by air or river travel and is located more than 1,000 km northeast of Lima. Conversely, NSI1428 was found almost a month later in Iquitos and has a different resistance pattern, suggesting this may be a different clone. However, more sequence analysis beyond the scope here would be required to assess the relatedness of these strains.

We report here the first instances of *bla*_{NDM-1} in *A. baumannii* in Peru, isolated from two different and geographically distinct regions at essentially the same time. This finding highlights the urgent need to implement surveillance and infection control measures for carbapenem-resistant pathogens, not only to mitigate the spread of resistant strains but also to limit or prevent the transfer of these resistance markers between pathogens in Peru and globally.¹

Received October 4, 2018. Accepted for publication December 3, 2018.

Published online January 21, 2019.

Disclosure: Claudio Rocha was an employee of the U.S. government during the conduct of this study. This work was prepared as part of his official duties. Title 17 U.S.C. § 105 provides that “Copyright protection under this title is not available for any work of the United States Government.” Title 17 U.S.C. § 101 defines a U.S. government work as a work prepared by a military service member or employee of the U.S. government as part of that person’s official duties. The views expressed in this article are those of the author and do not necessarily reflect the official policy or position of the Department of the Navy, Department of Defense, nor the U.S. government.

Authors’ addresses: Claudio Rocha, Manuela Bernal, Enrique Canal, Paul Rios, Rina Meza, Miguel Lopez, Rosa Burga, Ricardo Abadie, Melita Pizango, James Regeimbal, and Andrea McCoy, U. S. Naval Medical Research Unit No 6, Lima, Peru, E-mails: claudiourochacaldaron@gmail.com, manuela.m.bernal.in@mail.mil, enriquecanal@hotmail.com, paul.a.rios12.in@mail.mil, rina.a.meza2.fr@mail.mil, miguellogz2010@hotmail.com, rosaburga@gmail.com, reabadies@gmail.com, melepfilo@gmail.com, james.m.regeimbal.mil@mail.mil, and andrea.j.mccoy.mil@mail.mil. Elia Diaz and Alexander Briones, Hospital Regional de Loreto, Loreto, Peru, E-mails: elydid160@gmail.com and alexander.brionesalejos@gmail.com. Cesar Ramal-Asayag, Hospital Regional de Loreto, and Universidad Nacional de la Amazonia Peruana, Loreto, Peru, E-mail: ramalasayag@yahoo.fr. William Vicente, Instituto Nacional de Enfermedades Neoplásicas, Lima, Peru, E-mail: wvicente@blufsteinlab.com.

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

REFERENCES

1. World Health Organization, 2017. *Guidelines for the Prevention and Control of Carbapenem-Resistant Enterobacteriaceae, Acinetobacter baumannii and Pseudomonas aeruginosa in Health Care Facilities*. Geneva, Switzerland: WHO.
2. Dortet L, Poirel L, Nordmann P, 2014. Worldwide dissemination of the NDM-type carbapenemases in gram-negative bacteria. *Biomed Res Int* 2014: 249856.
3. Molton JS, Tambyah PA, Ang BS, Ling ML, Fisher DA, 2013. The global spread of healthcare-associated multidrug-resistant bacteria: a perspective from Asia. *Clin Infect Dis* 56: 1310–1318.
4. Stryko JP, Mony V, Cleveland J, Siddiqui H, Homel P, Gagliardo C, 2016. International travel is a risk factor for extended-spectrum β -lactamase-producing Enterobacteriaceae acquisition in children: a case-case-control study in an urban U.S. hospital. *Travel Med Infect Dis* 14: 568–571.
5. Seija V, Medina Presentado JC, Bado I, Papa Ezdra R, Batista N, Gutierrez C, Guirado M, Vidal M, Nin M, Vignoli R, 2015. Sepsis caused by New Delhi metallo- β -lactamase (*bla*_{NDM-1}) and qnrD-producing *Morganella morganii*, treated successfully with fosfomycin and meropenem: case report and literature review. *Int J Infect Dis* 30: 20–26.
6. Pilonetto M, Arend L, Vespero EC, Pelisson M, Chagas TP, Carvalho-Assef AP, Asensid MD, 2014. First report of NDM-1-producing *Acinetobacter baumannii* sequence type 25 in Brazil. *Antimicrob Agents Chemother* 58: 7592–7594.
7. Quiñones D et al., 2015. High prevalence of *bla* OXA-23 in *Acinetobacter* spp. and detection of *bla* NDM-1 in *A. soli* in Cuba: report from National Surveillance Program (2010–2012). *New Microbes New Infect* 7: 52–56.
8. Montaña S, Cittadini R, Del Castillo M, Uong S, Lazzaro T, Almuzara M, Barberis C, Vay C, Ramirez MS, 2016. Presence of New Delhi metallo- β -lactamase gene (*NDM-1*) in a clinical isolate of *Acinetobacter junii* in Argentina. *New Microbes New Infect* 11: 43–44.
9. Resurrección-Delgado C, Montenegro-Ildrogo JJ, Chiappe-Gonzalez A, Vargas-Gonzales R, Cucho-Espinoza C, Mamani-Condori DH, Huaroto-Valdivia LM, 2017. [Klebsiella pneumoniae NEW DELHI METALLO-LACTAMASE IN A PERUVIAN NATIONAL HOSPITAL]. *Rev Peru Med Exp Salud Publica* 34: 261–267.
10. Pendleton JN, Gorman SP, Gilmore BF, 2013. Clinical relevance of the ESKAPE pathogens. *Expert Rev Anti Infect Ther* 11: 297–308.
11. Clinical and Laboratory Standards Institute, 2017. *Performance Standards for Antimicrobial Susceptibility Testing; 27th Informational Supplement. CLSI M100-S27*. Wayne, PA: Clinical and Laboratory Standards Institute.
12. Magiorakos AP et al., 2012. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* 18: 268–281.
13. Poirel L, Walsh TR, Cuvillier V, Nordmann P, 2011. Multiplex PCR for detection of acquired carbapenemase genes. *Diagn Microbiol Infect Dis* 70: 119–123.
14. Ellington MJ, Kistler J, Livermore DM, Woodford N, 2007. Multiplex PCR for rapid detection of genes encoding acquired metallo- β -lactamases. *J Antimicrob Chemother* 59: 321–322.
15. Gonzalez-Villoria AM, Valverde-Garduno V, 2016. Antibiotic-resistant *Acinetobacter baumannii* increasing success remains a challenge as a nosocomial pathogen. *J Pathog* 2016: 7318075.
16. Peleg AY, Seifert H, Paterson DL, 2008. *Acinetobacter baumannii*: emergence of a successful pathogen. *Clin Microbiol Rev* 21: 538–582.
17. Higgins PG, Dammhayn C, Hackel M, Seifert H, 2010. Global spread of carbapenem-resistant *Acinetobacter baumannii*. *J Antimicrob Chemother* 65: 233–238.
18. Bonomo RA, Szabo D, 2006. Mechanisms of multidrug resistance in *Acinetobacter* species and *Pseudomonas aeruginosa*. *Clin Infect Dis* 43: S49–S56.
19. Bontron S, Nordmann P, Poirel L, 2016. Transposition of Tn125 encoding the NDM-1 carbapenemase in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 60: 7245–7251.
20. Bonnin RA, Poirel L, Naas T, Pirs M, Seme K, Schrenzel J, Nordmann P, 2012. Dissemination of New Delhi metallo- β -lactamase-1-producing *Acinetobacter baumannii* in Europe. *Clin Microbiol Infect* 18: E362–E365.